Lecture 15

Tuesday, 27 October 2026 8:17 PM

Tournament Solutions

We want to see when there is no Condorcet winner.

A voter $V \in \mathcal{N}$ has preferences $A \succ_V B \succ_V C \succ_V \dots$ (a strict total order) over all alternatives $A, B, C, \dots$ in $\mathcal{A}$.
The majority graph is a tournament.

$T = (V, A)$ is a tournament on vertices V.
Recall: Every tournament can be obtained as the majority graph of a profile.
Write the set of tournaments as a tournament, rather than a profile.
Notation: $a \to b$ means a beats b.

For a directed graph $G = (V, E)$, $u \rightsquigarrow v$ if there is a directed path from u to v. We assume $u \rightsquigarrow u$. Say u, v are strongly connected ($u \leftrightsquigarrow v$) if $u \rightsquigarrow v$ & $v \rightsquigarrow u$.

Claim: $\leftrightsquigarrow$ is an equivalence relation (reflexive, symmetric, transitive).

Def: $S \subseteq V$ is a strongly connected component if S is an equivalence class under $\leftrightsquigarrow$.

Consider G^*, obtained by replacing each component in G by a single vertex, and add edge (A, B) if there is an edge $a \to b$ in component A & b in component B.
G^* is called a condensation of G.


Claim: G^* is acyclic.

Hence G^* is a DAG, & hence the components in G can be written in topological order.

For a tournament, the condensation is a cyclic tournament.

A tournament rule is a partial choice correspondence $F: \mathcal{T} \to 2^{\mathcal{A}}$.
A tournament rule is Condorcet consistent if, whenever there is a Condorcet winner a, $F(T) = a$.

Consider: T_1:



Def: $X \subseteq A$ is dominant if $\forall x \to y$ $\forall a \in X$, $y \in X$.

Claim: If X, X' are two dominant sets, $X \neq X'$. Then either $X \subseteq X'$ or $X' \subseteq X$.
Proof: Suppose not. Then the fact $X \not\subseteq X'$, $x \in X \setminus X'$. But then since both X, X' are dominant, $x \to x'$ & $x' \to x$. ↯
Thus, dominant sets form a chain.
A top cycle is the minimal dominant, or a smallest dominant set (must exist, since A is a dominant set).

Claim: For a tournament, the components can be numbered $C_1, \dots, C_k$ s.t. $\forall i < j$, every vtx in C_i beats every vtx in C_j.
(easy)

Claim: The top cycle $TC(T) = C_1$.
(easy)

Claim: $TC(T)$ is Condorcet - consistent.
(easy)

Copeland: $cc(a) = |\{b : a \to b\}|$ = out-degree of a.
$(= \mu(a,b) > 0)$
Then $CO(T) = \{a : a$ has highest out-degree in $T\}$.

Claim: $CO(T) \subseteq TC(T)$
(easy)
Corollary: Copeland is Condorcet - consistent.

Given T, the dominion of a is $D(a) = \{b : a \to b\}$
Say b "loses" to $D(a) \setminus \{b\} \cup \{a\}$ & $D(a) \subseteq D(b)$
a is uncovered if no alternative covers a.
$UC(T)$ is the set of uncovered alternatives.

Claim: $UC(T) \neq \emptyset$
Proof: a) if $a \to b$ & $b \to c$, then $a \to c$
b) if $a \to b$ & $b \not\to a$
Then, this gives a strict partial order on the alternatives. Again, any acyclic, finite binary relation has undominated elements, & so $UC(T) \neq \emptyset$.

Def: $a \in A$ is a key if $\forall b \neq a$, either $a \to b$ or $\exists c : a \to c \to b$.

Claim: $a \in UC(T)$ iff a is a key.
Proof: Fix $a \in UC(T)$. WTS: $\forall b \neq a$, either $a \to b$ or $\exists c : a \to c \to b$.
Fix $b \neq a$, and suppose $a \not\to b$. Then $b \to a$.
Further, since $a \in UC$, $\exists c$ s.t. $a \to c$ & $c \not\to b$.
In the latter case, $b \to a$, & $a \to c$, so $b \to c$.
Now suppose a is a key. WTS $a \in UC(T)$.
Suppose for a contradiction $\exists b : b \to a$ and $a \to c \Rightarrow b \to c$.
Then a does not have a path to b of length ≤ 2, & so is not a key. ↯

Claim: $CO(T) \subseteq UC(T) \subseteq TC(T)$
Proof: a has max out-degree $\Rightarrow$ a is uncovered.
a is uncovered $\Rightarrow$ $a \in TC(T)$ (as $a \in C_1$)
(easy) ↯

So far, we've only looked at the unvoted tournament graph.
Suppose we also consider the "weighted" case.

Def: $\Delta(A) = \{x \in \mathbb{R}^n : \sum_{a \in A} x_a = 1\}$
$x \in \Delta(A)$ is a lottery over A (or a probability vector).

We know: every pair $a \neq b \in A$ s.t. $\forall b$, $\mu(a,b) \geq 0$
Does there exist a distribution p s.t. $\forall b$, $\mathop{\mathbb{E}}_{a \sim p}[\mu(a,b)] \geq 0$?

Define $M \in \mathbb{Z}^{n \times n}$: $M_{ab} = \mu(a,b)$
(the skew-symmetric extension of μ, & $M_{ba} = -M_{ab}$, or $M = -M^T$ & M is skew-symmetric)

Then for $p \in \Delta(A)$, $p^T M = \left(\sum_{a \in A} p_a \mu(a,b) \right)_{b \in A} = \left(\mathop{\mathbb{E}}_{a \sim p}[\mu(a,b)] \right)_{b \in A}$
& $p^T M_y = \sum_{a,b \in A} p_a y_b \mu(a,b)$

Def: A lottery p is maximal if $\forall y$, $p^T M y \geq 0$
$\Leftrightarrow$ $p^T M \geq 0$

Theorem: Maximal lotteries exist.
Proof: Consider:

max	λ		min	μ
$M^T p$	$\geq \lambda$	$\cdot q$	$-q^T M$	$\leq \mu$
$\sum p_a$	$= 1$	$\cdot \mu$	$\sum q_b$	$= 1$
p	≥ 0		q	≥ 0

Both feasible so WLOG have finite optimal values. say (p^*, λ^*), (q^*, μ^*)
Then $\lambda^* = \mu^*$.
Suppose $\lambda^* > 0$. Then OC $q^{*T} M^T p^* = -q^{*T} M p^*$
$\leq -\mu^* p^*$
< 0 ↯

Theorem: Let p be any maximal lottery.
① If a^* is a Condorcet winner then $p_{a^*} = 1$
② If $\hat{a}$ is a Condorcet loser then $p_{\hat{a}} = 0$

Proof: Fix ①: Let p be max. Then $\forall b$, $pM = \mu(a,b) > 0$
Hence p is maximal.
Now consider any lottery p, and $q : \mathbb{1}_{a^*}$.
Then $p^T M q = \sum_a p_a \mu(a, a^*) = \sum_{a \neq a^*} p_a \mu(a, a^*)$

If $\exists a \neq a^*$ s.t. $p_a > 0$, then this < 0. ↯
For ②

Claim: If p is a maximal lottery, then $\forall b$: $p_b > 0 \Rightarrow (pM)_b = 0$
Proof: Else $pM_b > 0$, but $pM p > 0$ for other p sometimes.

Proof of ②: $\forall p$, $(pM)_{\hat{a}} = \sum_{a \in A} p_a \mu(a, \hat{a})$. Now $\mu(a, \hat{a}) > 0$ used
$= \sum_{a \neq \hat{a}} p_a \mu(a, \hat{a})$

If $\exists a \neq \hat{a}$ s.t. $p_a > 0$ then $(p^T M)_{\hat{a}} > 0$
so: $(p^T M)_{\hat{a}} = 0 \Rightarrow p_a = 1$, but then p is not maximal.
So for any maximal lottery, $(pM)_{\hat{a}} > 0$
By the claim, this means $p_{\hat{a}} = 0$.

Example: 3: $a > b > c$
4: $b > c > a$
2: $c > a > b$



then: $M = \begin{array}{c} \\ a \\ b \\ c \end{array} \begin{array}{c} \begin{array}{ccc} a & b & c \end{array} \\ \begin{bmatrix} 0 & 1 & -5 \\ -1 & 0 & 3 \\ 5 & -3 & 0 \end{bmatrix} \end{array}$ $p = \begin{bmatrix} 5/9 \\ 3/9 \\ 1/9 \end{bmatrix}$